

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 July 2003 (17.07.2003)

PCT

(10) International Publication Number
WO 03/057220 A1

(51) International Patent Classification⁷: **A61K 31/4453**,
31/5355, 31/4166, 31/423, 31/44, C07D 233/70, 233/78,
413/04, 401/04, 491/08

Science Park South, Third Avenue, Harlow, Essex CM19
5AW (GB).

(21) International Application Number: PCT/GB03/00020

(74) Agent: **KONDO, Rie**; GlaxoSmithKline, CN925.1, 980
Great West Road, Brentford, Middlesex TW8 9GS (GB).

(22) International Filing Date: 7 January 2003 (07.01.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0200283.0 8 January 2002 (08.01.2002) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **GLAXO
GROUP LIMITED** [GB/GB]; Glaxo Wellcome House,
Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **BROMIDGE,
Steven, Mark** [GB/GB]; GlaxoSmithKline, New Frontiers
Science Park South, Third Avenue, Harlow, Essex CM19
5AW (GB). **LOVELL, Peter, John** [GB/GB]; Glaxo-
SmithKline, New Frontiers Science Park South, Third
Avenue, Harlow, Essex CM19 5AW (GB). **GOODACRE,
Caroline** [GB/GB]; GlaxoSmithKline, New Frontiers

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: CYCLIC UREA DERIVATIVES WITH 5-HT_{2C} RECEPTOR ACTIVITY




(57) Abstract: Compounds of formula (I) or a pharmaceutically acceptable salt thereof, having 5HT_{2C} receptor activity, are disclosed: wherein a is 0, 1, 2, 3, 4 or 5; b is 1, 2 or 3; Y is nitrogen or carbon; A is oxygen, nitrogen, -CONH-, -NHCO- or together with R² form a benzoxazolone group; R¹ is halogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, hydroxy, amino, mono- or di-C₁₋₆alkylamino, nitro, CN, CF₃, OCF₃, aryl, arylC₁₋₆alkyl, arylC₁₋₆alkyloxy or arylC₁₋₆alkylthio; R² is hydrogen, halogen, C₁₋₆alkyl, C₁₋₆alkoxy, haloC₁₋₆alkyl or haloC₁₋₆alkoxy; R³ is: (i) -NR⁴R⁵ where R⁴ and R⁵ are independently hydrogen, C₁₋₆alkyl or arylC₁₋₆alkyl; or (ii) an optionally substituted N-linked heterocycle; or (iii) an optionally substituted C-linked heterocycle; ===== is a single bond or a double bond; and X is CH₂ or C=O (when ===== is a single bond) or X is CH (when ===== is a double bond). The preparation of these compounds and their use in therapy, especially for a CNS disorder such as depression or anxiety, are also disclosed.

Cyclic Urea Derivatives Possessing 5-HT_{2C} Receptor Activity,
Preparation Thereof And Use In Therapy

This invention relates to novel compounds having pharmacological activity, processes for
 5 their preparation, to compositions containing them and to their use in the treatment and/or
 prevention of CNS and other disorders.

WO 96/23783, WO 97/46699 and WO 97/48700 all disclose a series of indoline derivatives
 10 which are 5-HT_{2C} receptor antagonists and which are claimed to be useful in the treatment of
 various CNS disorders.

A novel class of compounds possessing 5-HT_{2C} receptor activity has been found. The
 present invention therefore provides, in a first aspect, a compound of formula (I) or a
 15 pharmaceutically acceptable salt thereof:



(I)

wherein:

a is 0, 1, 2, 3, 4 or 5;

b is 1, 2 or 3;

20 Y is nitrogen or carbon;

A is oxygen, nitrogen, -CONH-, -NHCO- or together with R² form a benzoxazolone group;

R¹ is halogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, hydroxy, amino, mono- or
 di-C₁₋₆alkylamino, nitro, CN, CF₃, OCF₃, aryl, arylC₁₋₆alkyl, arylC₁₋₆alkyloxy or
 arylC₁₋₆alkylthio;

25 R² is hydrogen, halogen, C₁₋₆alkyl, C₁₋₆alkoxy, haloC₁₋₆alkyl or haloC₁₋₆alkoxy;

R³ is: (i) -NR⁴R⁵ where R⁴ and R⁵ are independently hydrogen, C₁₋₆alkyl or arylC₁₋₆alkyl; or

(ii) an optionally substituted N-linked heterocycle; or

(iii) an optionally substituted C-linked heterocycle;

30 ----- is a single bond or a double bond;

X is CH₂ or C=O (when ----- is a single bond) or X is CH (when ----- is a double bond).

The following terms, whether used alone or as part of another group are to be given the
 35 following meanings, unless otherwise stated.

The term "halogen" and its abbreviated form "halo" are used herein to describe fluorine, chlorine, bromine or iodine.

5 The term "alkyl" is used herein to describe a straight chain or branched fully saturated hydrocarbon group. "C₁₋₆alkyl" refers to alkyl groups having from one to six carbon atoms, including all isomeric forms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, neopentyl, sec-pentyl, n-pentyl, isopentyl, tert-pentyl and hexyl.

10 The term "C₁₋₆alkoxy" refers to a straight chain or branched chain alkoxy (or "alkyloxy") group having from one to six carbon atoms, including all isomeric forms, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentoxy, neopentoxy, sec-pentoxy, n-pentoxy, isopentoxy, tert-pentoxy and hexoxy.

15 The term "C₁₋₆alkylthio" refers to a straight chain or branched chain alkylthio group having from one to six carbon atoms, such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, tert-butylthio, pentylthio, neopentylthio, sec-pentylthio, n-pentylthio, isopentylthio, tert-pentylthio and hexylthio.

20 The term "mono- or di-C₁₋₆alkylamino" refers to an amino group which is substituted by one C₁₋₆alkyl group or an amino group which is substituted by two C₁₋₆alkyl groups, the two amino groups being the same or different. Examples of mono-C₁₋₆alkylamino include methylamine, ethylamine, propylamine, isopropylamine, butylamine, isobutylamine, sec-butylamine, tert-butylamine, pentylamine, neopentylamine, sec-pentylamine, n-pentylamine, isopentylamine, tert-pentylamine and hexylamine. Examples of di-C₁₋₆alkylamino include
25 dimethylamine, diethylamine, dipropylamine, diisopropylamine, dibutylamine, diisobutylamine, disec-butylamine, ditert-butylamine, dipentylamine, dineopentylamine, dihexylamine, butylmethylamino, isopropylmethylamino, ethylisopropylamino, ethylmethylamino, etc.

30 The term "aryl" is used herein to describe groups such as phenyl or naphthyl, which may be optionally substituted by one or more of C₁₋₆alkyl (to form "arylC₁₋₆alkyl"), halogen, CF₃ or C₁₋₆alkoxy (to form "arylC₁₋₆alkoxy").

35 The terms "halo C₁₋₆alkoxy" or "haloC₁₋₆alkyl" are used to describe a C₁₋₆alkoxy or a C₁₋₆alkyl group, respectively, substituted with one or more halogens. Examples include -CHCl₂, -CF₃, -OCF₃, etc.

40 The term "optionally substituted N-linked heterocycle" is used herein to describe a stable non-aromatic 5-7 membered ring containing at least 1 nitrogen atom and optionally a further 1 or 2 heteroatoms selected from nitrogen, sulphur or oxygen, wherein the heterocycle is linked to the remainder of the molecule via a nitrogen atom.

The term "optionally substituted C-linked heterocycle" is used herein to describe a stable non-aromatic 5-7 membered ring containing at least 1 nitrogen atom and optionally a further 1 or 2 heteroatoms selected from nitrogen, sulphur and oxygen, wherein the heterocycle is linked to the remainder of the molecule via a carbon atom.

5

Suitable examples of N-linked or C-linked heterocycles include pyrrolidinyl, piperazinyl, morpholinyl, imidazolidinyl, thiomorpholinyl, piperidinyl and azepanyl.

More than one optional substituent may be present in the N-linked or C-linked heterocycle, which may be the same or different, and may be attached to any carbon atom of the heterocycle or an available nitrogen atom.

10

Suitable optional substituents for the N-linked or C-linked heterocycle include C₁₋₆alkyl, amino, mono- or di- C₁₋₆alkylamino, aryl, arylC₁₋₆alkyl, arylamino, hydroxy, C₁₋₆alkylamido, hydroxyC₁₋₆alkyl, C₁₋₆alkoxycarbonyl, halogen, haloC₁₋₆alkyl, a heteroaromatic group (such as indole or benzimidazole), an aromatic or non-aromatic N-linked or C-linked heterocycle or an aromatic or non-aromatic heterocycleC₁₋₆alkyl optionally substituted by C₁₋₆alkyl. Examples of aromatic or non-aromatic heterocycleC₁₋₆alkyl include heterocycle-methyl (such as pyridinyl-methyl and benzimidazolyl-methyl) and heterocycle-ethyl (such as morpholinyl-ethyl and indolyl-ethyl).

15


20

Substituents in the N-linked or C-linked heterocycle may form a bridge structure, to form a group such as for example 2-oxa-5-azabicyclo[2.2.1]heptyl. Such a bicyclic group may be further substituted by the substituents listed above. More than one substituent may be present on the same carbon atom to form spiro structures such as 1,4 and 1,5 dioxo spiro compounds.

25

When a is 2 or more, the 2 or more R¹ groups may be the same or different.

When A is nitrogen, A and R² may be joined to form a benzoxazolone group as follows:



30

Preferred compounds of the present invention include those in which, independently or in any compatible combination:

a is 1 or 2;

R¹ is halogen, particularly fluoro or chloro;


R¹ is at the 3 or 4 position, ie at the *meta* and/or *para* positions of the benzene to which R¹ is attached;

35

X is CH₂ and ----- represents a single bond, or X is CH and ----- represents a double bond;

Y is carbon;


R² is at the following position:



5

R² is C₁₋₆alkoxy, particularly methoxy;

A is at the following position:



10

A is oxygen;

b is 2;

R³ is an N-linked heterocycle, particularly piperidinyl, or is a diC₁₋₆alkylamine, particularly dimethylamine.

15 Exemplary compounds of this invention include:

1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one

1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one

1-(4-Methyl-3-trifluoromethyl-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one

20

1-(4-Methyl-3-trifluoromethyl-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one

1-(2-Chloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one

1-[4-Methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-3-phenyl-imidazolidin-2-one

1-[3-(2-Dimethylamino-ethoxy)-4-methoxy-phenyl]-3-(3-fluorophenyl)-imidazolidin-2-one

25

1-(3-Fluoro-phenyl)-3-{4-methoxy-3[(1S,4S)-2-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-ethoxy]-phenyl}-imidazolidin-2-one

1-(3,4-Dichloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one

30

1-(3,4-Dichloro-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one

1-(3,4-Dichloro-phenyl)-3-[3-(2-dimethylamino-ethoxy)-4-methoxy-phenyl]-imidazolidin-2-one

1-(3,4-Dichloro-phenyl)-3-[6-(2-piperidin-1-yl-ethoxy)-pyridin-3-yl]-imidazolidin-2-one
 1-[4-Bromo-3-(2-piperidin-1-yl-ethoxy)-phenyl]-3-(3,4-dichloro-phenyl)-imidazolidin-2-one
 1-(2,3-Dichloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one

- 5 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-1,3-dihydro-imidazol-2-one
 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-1,3-dihydro-imidazol-2-one
 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazol-2-one
 10 3-(3-Fluoro-phenyl)-1-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazol-2,4-dione
 5-[3-(3,4-Dichloro-phenyl)-2-oxo-imidazolidin-1-yl]-3-(2-piperidin-1-yl-ethyl)-3H-benzoxazol-2-one
 and pharmaceutically acceptable salts thereof.


- 15 The compounds of formula (I) can form acid addition salts. It will be appreciated that for use in medicine the salts of the compounds of formula (I) should be pharmaceutically acceptable. Suitable pharmaceutically acceptable salts will be apparent to those skilled in the art and include those described in J. Pharm. Sci., 1977, 66, 1-19, such as acid addition salts formed with inorganic acids e.g. hydrochloric, hydrobromic, sulfuric, nitric or phosphoric acid; and
 20 organic acids e.g. succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, p-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid.

The compounds of this invention may be in crystalline or non-crystalline form, and, if crystalline, may optionally be hydrated or solvated. This invention includes within its scope
 25 stoichiometric hydrates as well as compounds containing variable amounts of water.

- Certain compounds of formula (I) are capable of existing in stereoisomeric forms (e.g. geometric or ("cis-trans") isomers, diastereomers and enantiomers) and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates.
 30 The different stereoisomeric forms may be separated one from the other by the usual methods, or any given isomer may be obtained by stereospecific or asymmetric synthesis. The invention also extends to any tautomeric forms and mixtures thereof.

- The present invention also provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises:
 35


a) the cyclisation of a compound of formula (II)



in which R^1 , R^2 , R^3 , A, Y, a and b are defined in formula (I); or


b) the coupling of a compound of formula (III):

5



(III)


10 in which R^1 , R^2 , A, X, Y, a and b are defined in formula (I) and L^1 is a leaving group, with a compound of formula (IV):



(IV)


15 in which R^4 and R^5 are defined in formula (I); or

c) the coupling of a compound of formula (V)



(V)


20 in which R^1 , R^2 , X, Y and a are defined in formula (I), with a compound of formula (VI):



(VI)

25 in which R^4 , R^5 and b are defined in formula (I) and L^2 is a leaving group or hydroxy; or

d) the cyclisation of a compound of formula (VII)



(VII)

in which R^1 , R^2 , R^3 , Y , A , a and b are defined in formula (I);

optionally followed by:

- removing any protecting groups; and/or
- converting a compound of formula (I) into another compound of formula (I); and/or
- forming a pharmaceutically acceptable salt.

For process a), suitable cyclising agents include phosgene, triphosgene and CDI (preferably phosgene). The cyclisation reaction of compounds of formula (II) is preferably carried out in an inert solvent such as tetrahydrofuran in the presence of a base such as triethylamine.

For process b), suitably L^1 is mesylate, tosylate, bromo or chloro (preferably mesylate). The reaction of compounds of formulae (III) and (IV) are typically carried out in the presence of a base such as potassium carbonate or sodium carbonate (preferably potassium carbonate) in a suitable solvent such as N,N-dimethylformamide.


For process c), suitably L^2 is halogen (preferably chloro) or hydroxy. When L^2 is halogen, the reaction of compounds of formulae (V) and (VI) are typically carried out in the presence of a base such as potassium carbonate or sodium carbonate (preferably potassium carbonate) in a suitable solvent such as ethylene glycol dimethyl ether. When L^2 is hydroxy, suitable Mitsunobu reagents include triphenylphosphine and DEAD in a suitable solvent such as tetrahydrofuran.

For process d), suitable cyclising conditions include catalytic acid (preferably HCl) in an inert solvent such as toluene.


Compounds of formula (I) can be converted into further compounds of formula (I) using standard techniques. For example, and by way of illustration rather than limitation, compounds of formula (I) in which $----$ is a double bond can be converted to compounds of formula (I) in which $----$ is a single bond by palladium catalysed hydrogenation in a suitable solvent such as ethanol.

Compounds of formulae (IV) and (VI) are commercially available. Compounds of formulae (II), (III), (V) and (VII) may be prepared according to methods described herein, or may be prepared according to known methods or by analogous methods thereto.

For example, a compound of formula (II) may be prepared by reacting a compound of formula (VIII):




with a compound of formula (IX):




wherein L_3 is a leaving group.

A compound of formula (III) may be prepared by reacting a compound of formula (V) with a compound of formula (X):




(X) wherein L^4 is a leaving group, followed by subsequent debenzylation and activation of the alcohol to a leaving group (L^1).

A compound of formula (V) may be prepared by the phosgene cyclisation of a compound of formula (XI)



followed by subsequent debenzylation.

A compound of formula (VII) may be prepared for example by reacting a compound of formula (XII):



with phenyl isocyanate having the desired $[R^1]_n$ substituents. Compounds of formula (XII) may be prepared for example by reacting a compound of formula (VIII) with glyoxal 1,1-dimethyl acetal under reducing conditions, such as catalytic hydrogenation over a palladium catalyst.

Those skilled in the art will appreciate that it may be necessary to protect certain groups to carry out the above processes. Suitable protecting groups and methods for their attachment and removal are conventional in the art of organic chemistry, such as those described in Greene T.W. 'Protective groups in organic synthesis' New York, Wiley (1981).

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

In another aspect, the present invention provides a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

In a further aspect, the present invention provides a process for preparing a pharmaceutical composition, the process comprising mixing a compound of formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier or excipient.

A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusible solutions or suspensions or suppositories. Orally administrable compositions are generally preferred.

Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); tableting lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycollate); and acceptable wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated according to methods well known in normal pharmaceutical practice.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents (e.g. sorbitol syrup; cellulose derivatives or hydrogenated edible fats), emulsifying agents (e.g. lecithin or acacia), non-aqueous vehicles (which may include edible oils e.g. almond oil, oily esters, ethyl alcohol or fractionated vegetable oils), preservatives (e.g. methyl or propyl-p-hydroxybenzoates or sorbic acid), and, if desired, conventional flavourings or colorants, buffer salts and sweetening agents as appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. Formulations for injection may be presented in unit dosage form e.g. in ampoules or in multi-dose, utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle, optionally with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use. The compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilisation cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents. Drops may be formulated with an aqueous or non-aqueous base also comprising one or more dispersing agents, stabilising agents, solubilising agents or suspending agents. They may also contain a preservative.

The compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

The compounds of the invention may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or

intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

5 For intranasal administration, the compounds of the invention may be formulated as solutions for administration via a suitable metered or unitary dose device or alternatively as a powder mix with a suitable carrier for administration using a suitable delivery device. Thus compounds of formula (I) may be formulated for oral, buccal, parenteral, topical (including
10 ophthalmic and nasal), depot or rectal administration or in a form suitable for administration by inhalation or insufflation (either through the mouth or nose).

The compounds of the invention may be formulated for topical administration in the form of ointments, creams, gels, lotions, pessaries, aerosols or drops (e.g. eye, ear or nose drops).
15 Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Ointments for administration to the eye may be manufactured in a sterile manner using sterilised components.

Compounds of formula (I) and their pharmaceutically acceptable salts are of use in the
20 treatment of certain CNS disorders such as depression, which term is used herein to include bipolar depression, unipolar depression, single or recurrent major depressive episodes with or without psychotic features, catatonic features, melancholic features, atypical features or postpartum onset, seasonal affective disorder, dysthymic disorders with early or late onset and with or without atypical features, neurotic depression and social phobia, depression
25 accompanying dementia for example of the Alzheimer's type, vascular dementia with depressed mood, schizoaffective disorder or the depressed type, and depressive disorders resulting from general medical conditions including, but not limited to, myocardial infarction, diabetes, miscarriage or abortion, *etc.* Other CNS disorders which may be treated or prevented include anxiety disorders, including generalised anxiety, schizophrenia, panic
30 disorder, agoraphobia, social phobia, obsessive compulsive disorder and post-traumatic stress disorder, pain (particularly neuropathic pain), memory disorders, including dementia, amnesic disorders and age-associated memory impairment, disorders of eating behaviours, including anorexia nervosa and bulimia nervosa, sexual dysfunction, sleep disorders (including disturbances of circadian rhythm, dyssomnia, insomnia, sleep apnea and narcolepsy),
35 withdrawal from abuse of drugs such as of cocaine, ethanol, nicotine, benzodiazepines, alcohol, caffeine, phencyclidine (phencyclidine-like compounds), opiates (e.g. cannabis, heroin, morphine), sedative ipnotic, amphetamine or amphetamine-related drugs (e.g. dextroamphetamine, methylamphetamine) or a combination thereof, motor disorders such as Parkinson's disease, dementia in Parkinson's disease, neuroleptic-induced Parkinsonism and
40 tardive dyskinesias, as well as other psychiatric disorders.

Compounds of formula (I) may also have utility in the treatment of certain gastrointestinal disorders such as irritable bowel syndrome, Crohn's disease, ulcerative colitis, non-steroidal anti-inflammatory drug induced damage.

- 5 As used herein, the term "treatment" refers to amelioration and/or cure of established symptoms as well as prophylaxis.

Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance, in particular in the treatment of the above
10 disorders. In particular the invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use as a therapeutic substance in the treatment of depression.

Compounds of the invention may be administered in combination with other active
15 substances such as 5HT3 antagonists, NK-1 antagonists, serotonin agonists, selective serotonin reuptake inhibitors (SSRI), noradrenaline re-uptake inhibitors (SNRI), tricyclic antidepressants and/or dopaminergic antidepressants.

Suitable 5HT3 antagonists which may be used in combination of the compounds of the
20 inventions include for example ondansetron, granisetron, metoclopramide.

Suitable serotonin agonists which may be used in combination with the compounds of the invention include sumatriptan, rauwolscine, yohimbine, metoclopramide.

25 Suitable SSRIs which may be used in combination with the compounds of the invention include fluoxetine, citalopram, femoxetine, fluvoxamine, paroxetine, indalpine, sertraline, zimeldine.

Suitable SNRIs which may be used in combination with the compounds of the invention
30 include venlafaxine and reboxetine.

Suitable tricyclic antidepressants which may be used in combination with a compound of the invention include imipramine, amitriptyline, chlomipramine and nortriptyline.

35 Suitable dopaminergic antidepressants which may be used in combination with a compound of the invention include bupropion and amineptine.

It will be appreciated that the compounds of the combination or composition may be administered simultaneously (either in the same or different pharmaceutical formulations),
40 separately or sequentially.

The invention further provides a method of treatment of the above disorders in mammals including humans, which comprises administering to the sufferer a therapeutically safe and effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

- 5 In another aspect, the invention provides for the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the treatment of the above disorders.

- 10 The composition of the present invention may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration. The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 1.0 to 200 mg, and such unit doses may be administered more
15 than once a day, for example two or three times a day. Such therapy may extend for a number of weeks or months. When administered in accordance with the invention, no unacceptable toxicological effects are expected with the compounds of the invention.

- 20 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

- 25 The following Descriptions and Examples illustrate the preparation of compounds of the present invention.

Description 1

2-(3-Fluoro-phenylamino)-ethanol (D1)

- 30 To a solution of 3-fluoroaniline (5 g, 45 mmol) in dichloromethane (100 mL) and pyridine (4.6 mL, 57 mmol) at room temperature was slowly added 2-chloroethylchloroformate (4.6 mL, 45 mmol). The mixture was stirred at room temperature for 2.5 hours and washed with water (x4), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was dissolved in ethanol (100 mL), treated with potassium hydroxide (10.1 g, 180
35 mmol) and heated at 90°C for 4 hours. The mixture was concentrated *in vacuo* and the residue partitioned between dichloromethane (250 mL) and water (150 mL). The aqueous layer was re-extracted and the combined organics washed with water (x5) and brine. The organics were dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as an orange oil
40 (5.6 g, 81%). MH^+ 156.

Description 2

2-(4-Methyl-3-trifluoromethyl-phenylamino)-ethanol (D2)

The title compound was prepared from 5-amino-2-methylbenzotrifluoride using the method described in Description 1. MH^+ 220.

5 **Description 3**

2-(2-Chloro-phenylamino)-ethanol (D3)

The title compound was prepared from 2-chloroaniline using the method described in Description 1. $(M-H_2O + H)^+$ 154/156.

10

Description 4

2-(3,4-Dichloro-phenylamino)-ethanol (D4)

The title compound was prepared from 3,4-dichloroaniline using the method described in Description 1. MH^+ 206/208/210.

15

Description 5

2-(2,3-Dichloro-phenylamino)-ethanol (D5)

20 The title compound was prepared from 2,3-dichloroaniline using the method described in Description 1. MH^+ 206/208/210.

Description 6

2-Benzyloxy-1-methoxy-4-nitro-benzene (D6)

25

A mixture of benzyl bromide (3.1 mL, 26.6 mmol), 2-methoxy-5-nitrophenol (3 g, 17.7 mmol) and sodium iodide (0.07 g, 0.5 mmol) in ethylene glycol dimethyl ether (100 mL), water (10 mL) and saturated potassium carbonate solution (35 mL) was heated at 65°C for 23 hours. The mixture was concentrated *in vacuo* and the residue partitioned between ethyl acetate (200 mL) and saturated sodium bicarbonate solution (75 mL). The organic phase was washed with brine (125 mL, 100 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a yellow solid (4.5 g, 98%).

30

35 **Description 7**

3-Benzyloxy-4-methoxy-phenylamine (D7)

Iron powder (0.85 g, 15.2 mmol) was added to a suspension of D6 (1.7 g, 6.6 mmol) in methanol (75 mL) and saturated aqueous ammonium chloride (35 mL) and heated at reflux for 2 hours. The mixture was allowed to cool to room temperature and filtered through a plug of celite and washed with methanol. The methanol was removed *in vacuo* and the residue partitioned between ethyl acetate (125 mL) and water (enough to dissolve the ammonium chloride). The aqueous layer was re-extracted with ethyl acetate (50 mL) and the combined

40

organics washed with brine (125 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a brown solid (1.24 g, 83%). MH^+ 230.

5 **Description 8**

1-[2-(2-Methoxy-5-nitro-phenoxy)-ethyl]-piperidine (D8)

10 A mixture of 2-methoxy-5-nitrophenol (15 g, 88.8 mmol) and 1-(2-chloroethyl)piperidine monohydrochloride (21.2 g, 115 mmol) in ethylene glycol dimethyl ether (600 mL), water (226 mL) and saturated potassium carbonate solution (226 mL) was stirred at room temperature for 24 hours. The organic phase was separated, washed with water (400 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo* to give the title compound as brown crystals (24.6 g, 98%). MH^+ 281.

15 **Description 9**

4-Methoxy-3-(2-piperidin-1-yl-ethoxy)-phenylamine (D9)

20 D8 (24.6 g, 87.9 mmol) was dissolved in ethanol (500 mL). 10% Palladium on charcoal (2 g) was added and the reaction mixture hydrogenated at atmospheric pressure for 16 hours. The mixture was filtered through a small plug of celite and the solvent removed *in vacuo* to give the title compound as brown crystals (21.8 g, 100%). MH^+ 251.

Description 10

4-[2-(2-Methoxy-5-nitro-phenoxy)-ethyl]-morpholine (D10)

25

The title compound was prepared from 2-methoxy-5-nitrophenol and 4-(2-chloroethyl)morpholine monohydrochloride stirring for 48 hours using the method described in Description 8. MH^+ 283.

30 **Description 11**

4-Methoxy-3-(2-morpholin-4-yl-ethoxy)-phenylamine (D11)

The title compound was prepared from D10 using the method described in Description 9. MH^+ 253.

35

Description 12

[2-(2-Methoxy-5-nitro-phenoxy)-ethyl]-dimethyl-amine (D12)

40 Diethyl azodicarboxylate (4.6 mL, 29 mmol) was added dropwise to a stirred solution of 2-methoxy-5-nitrophenol (4.97 g, 29 mmol), triphenylphosphine (7.7 g, 29 mmol) and N,N-dimethylethanolamine (2.95 mL, 29 mmol) at room temperature. After stirring for 48 hours the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (200 mL) and saturated sodium bicarbonate solution (125 mL). The organic phase was washed with

brine (2x100 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a yellow solid (1.99 g, 28%). MH^+ 241.

5 **Description 13**

3-(2-Dimethylamino-ethoxy)-4-methoxy-phenylamine (D13)

The title compound was prepared from D12 using the method described in Description 9.

10 **Description 14**

5-Nitro-2-(2-piperidin-1-yl-ethoxy)-pyridine (D14)

15 N-(2-Hydroxyethyl) piperidine (2.44 g, 18.9 mmol) was dissolved in N,N-dimethylformamide (40 mL), cooled to 0°C in an ice/water bath and sodium hydride (60% dispersion in oil, 0.682 g, 17 mmol) added portionwise over ten minutes. After stirring for 35 minutes at 0°C the reaction mixture was allowed to warm to room temperature and was stirred for 2 hours. The mixture was cooled to 0°C and 2-chloro-5-nitropyridine (3.0 g, 18.9 mmol) added portionwise. The mixture was allowed warm to room temperature and stirred for 16 hours. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate 20 (150 mL) and saturated sodium bicarbonate solution (250 mL). The organic phase was washed with brine (150 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a brown solid (0.897 g, 19%). MH^+ 252.

25 **Description 15**

6-(2-Piperidin-1-yl-ethoxy)-pyridin-3-ylamine (D15)

30 D14 (0.435 g, 1.7 mmol) was dissolved in ethanol (50 mL). 10% Palladium on charcoal (0.15 g) was added and the reaction mixture hydrogenated at atmospheric pressure for 2 hours and 20 minutes. The mixture was filtered through a small plug of celite and the solvent removed *in vacuo* to give the title compound as a grey solid (0.373 g, 97%).

Description 16

N-(4-Bromo-3-hydroxy-phenyl)-acetamide (D16)

35

3-Acetamidophenol (45.5 g, 0.3 mol) in acetic acid (400 mL) was treated with bromine (15.5 ml, 0.3 mol) in acetic acid (100 mL) at room temperature with stirring. The mixture was stirred at room temperature for 60 hours and filtered to give the title product as a white solid (67.55 g, 97 %). MH^+ 228/230.

40

Description 17

N-[4-Bromo-3-(2-piperidin-1-yl-ethoxy)-phenyl]-acetamide (D17)

A mixture of D16 (4.6 g, 20 mmol) and 1-(2-chloroethyl)piperidine monohydrochloride (4.05 g, 22 mmol) in ethylene glycol dimethyl ether (200 mL), water (50 mL) and saturated potassium carbonate solution (50 mL) was stirred at 70°C for three hours and at room temperature for 16 hours. The organic phase was separated, the aqueous layer extracted with dichloromethane, and the combined organics dried over anhydrous magnesium sulphate. The solvent was removed *in vacuo* to give the title compound as a white solid (4.2 g, 62%). MH^+ 341/343.

10 **Description 18**

4-Bromo-3-(2-piperidin-1-yl)-ethoxy)-phenylamine (D18)

A solution of D17 (2 g, 5.9 mmol) in ethanol (20 mL) and 40% sodium hydroxide solution (20 mL) was heated at 80°C for two hours. The reaction mixture was cooled and the organic layer separated and concentrated *in vacuo*. The residue was partitioned between dichloromethane and water and the organic layer dried over anhydrous magnesium sulphate. The solvent was removed *in vacuo* to give the title compound as a yellow solid (1.7 g, 97%). MH^+ 298/300

20 **Description 19**

(2,2-Dimethoxy-ethyl)-[4-methoxy-3-(2-piperidin-1-yl)-ethoxy)-phenyl]-amine (D19)

To a solution of D9 (2.49 g, 10 mmol) and glyoxal 1,1-dimethyl acetal solution (~45% in tert-butyl methyl ether, 2.8 mL, 12 mmol) in ethanol (50 mL) was added 10% palladium/carbon (0.45 g) and the mixture hydrogenated at atmospheric pressure for 31 hours. The mixture was filtered through a small plug of celite and the solvent removed *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a brown oil (1.81 g, 54%). MH^+ 339.

30 **Description 20**

1-(2,2-Dimethoxy-ethyl)-3-(3-fluoro-phenyl)-1-[4-methoxy-3-(2-piperidin-1-yl)-ethoxy)-phenyl]-urea (D20)

3-Fluorophenyl isocyanate (0.186 g, 1.4 mmol) in dichloromethane (1 mL) was added to a solution of D19 (0.455 g, 1.3 mmol) in dichloromethane (15 mL) at room temperature. After stirring for 16 hours at room temperature the reaction mixture was diluted with dichloromethane (45 mL), washed with saturated sodium bicarbonate solution (20 mL) and brine (50 mL). The organic phase was dried over anhydrous magnesium sulphate and the solvent removed *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a white solid (0.316 g, 49%). MH^+ 476.

Description 21

3-(3,5-Difluoro-phenyl)-1-(2,2-dimethoxy-ethyl)-1-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-urea

The title compound was prepared from 3,5-difluorophenyl isocyanate and D19 using the method described in Description 20. MH⁺ 494.

Description 22

(3-Benzoyloxy-4-methoxy-phenylamino)-acetic acid ethyl ester (D22)

D7 (1.5 g, 6.6 mmol) was dissolved in N,N-dimethylformamide (20 mL) and treated with potassium carbonate (1.81 g, 13.1 mmol), ethyl bromoacetate (0.73 mL, 6.6 mmol) and sodium iodide (0.983 g, 6.6 mmol). The reaction mixture was heated at 80°C for 17 hours and allowed to stand at room temperature for 48 hours. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (125 mL) and half saturated sodium bicarbonate solution (85 mL). The organic phase was washed with brine (150 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound (1.3 g, 63%). MH⁺ 316.

Description 23

1-(3-Benzoyloxy-4-methoxy-phenyl)-3-(3-fluoro-phenyl)-imidazolidine-2,4-dione (D23)

D22 (1.3 g, 4.1 mmol) was dissolved in toluene (50 mL) and 3-fluorophenyl isocyanate (0.566 g, 4.1 mmol) added. The reaction mixture was heated at 85°C for ten hours. The solvent was removed *in vacuo* and the residue purified by silica gel chromatography to give the title compound as a pale yellow solid (1.03 g, 61%). MH⁺ 407.

Description 24

N-(3-Benzoyloxy-4-methoxy-phenyl)-N'-(3-fluoro-phenyl)-ethane-1,2-diamine (D24)

Methanesulfonyl chloride (0.68 mL, 8.8 mmol) was added dropwise to a solution of D1 (1.36 g, 8.8 mmol) in dichloromethane (17 mL) and triethylamine (1.83 mL, 13.2 mmol) at room temperature and stirred for 30 minutes. The mixture was diluted with dichloromethane (120 mL) and washed with water (75 mL) and brine (115 mL). The organic phase was dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue and sodium iodide (1.45 g, 9.7 mmol) were added at 50°C to a mixture of D7 (2.3 g, 10 mmol) and potassium carbonate (1.22 g, 8.8 mmol) in N,N-dimethylformamide (17 mL). The reaction mixture was heated at 80°C for 16 hours and allowed to cool to room temperature. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (160 mL) and water (100 mL). The organic phase was washed with brine (175 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a brown oil (1.5 g, 47%). MH⁺ 367.

Description 25

N-(3-Benzoyloxy-4-methoxy-phenyl)-N'-(4-methyl-3-trifluoromethyl-phenyl)-ethane-1,2-diamine (D25)

5 The title compound was prepared from D2 and D7 using the method described in Description 24. MH^+ 431.

Description 26

N-(3-Benzoyloxy-4-methoxy-phenyl)-N'-(2-chloro-phenyl)-ethane-1,2-diamine (D26)

10 The title compound was prepared from D3 and D7 using the method described in Description 24. MH^+ 382/384

Description 27

15 **N-(3,4-Dichloro-phenyl)-N'-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-ethane-1,2-diamine (D27)**

The title compound was prepared from D4 and D9 using the method described in Description 24. MH^+ 438/440/442.

20 **Description 28**

N-(3,4-Dichloro-phenyl)-N'-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-ethane-1,2-diamine (D28)

25 The title compound was prepared from D4 and D11 using the method described in Description 24. MH^+ 440/442/444.

Description 29

30 **N-(3,4-Dichloro-phenyl)-N'-[3-(2-dimethylamino-ethoxy)-4-methoxy-phenyl]-ethane-1,2-diamine (D29)**

The title compound was prepared from D4 and D13 using the method described in Description 24. MH^+ 398/400/402.

Description 30

35 **N-(3,4-Dichloro-phenyl)-N'-[6-(2-piperidinyl-1-yl-ethoxy)-pyridin-3-yl]-ethane-1,2-diamine (D30)**

The title compound was prepared from D4 and D15 using the method described in Description 24. MH^+ 409/411/413.

40

Description 31

N-[4-Bromo-3-(2-piperidin-1-yl-ethoxy)-phenyl]-N'-(3,4-dichloro-phenyl)-ethane-1,2-diamine (D31)

The title compound was prepared from D4 and D18 using the method described in Description 24. MH^+ 485/487/489/491

5 **Description 32**

N-(2,3-Dichloro-phenyl)-N'-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-ethane-1,2-diamine (D32)

10 The title compound was prepared from D5 and D9 using the method described in Description 24. MH^+ 438/440/442

Description 33

1-(3-Benzyloxy-4-methoxy-phenyl)-3-(3-fluoro-phenyl)-imidazolidin-2-one (D33)

15 Phosgene solution (~20% in toluene, 2.1 mL, 4.3 mmol) was added dropwise at room temperature to a solution of D24 (1.48 g, 4.0 mmol) in tetrahydrofuran (55 mL) and triethylamine (1.7 mL 12 mmol). After stirring for 2.5 hours the solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and 2M sodium hydroxide solution. The organic phase was washed with brine, dried over anhydrous magnesium sulphate and
20 concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a white solid (1.16 g, 73%). MH^+ 414.

Description 34

25 **1-(3-Benzyloxy-4-methoxy-phenyl)-3-(4-methyl-3-trifluoromethyl-phenyl)-imidazolidin-2-one (D34)**

The title compound was prepared from D25 using the method described in D33. MH^+ 457.

Description 35

30 **1-(3-Benzyloxy-4-methoxy-phenyl)-3-(2-chloro-phenyl)-imidazolidin-2-one (D35)**

The title compound was prepared from D26 using the method described in D33. MH^+ 430/432.

35 **Description 36**

1-(3-Fluoro-phenyl)-3-(3-hydroxy-4-methoxy-phenyl)-imidazolidin-2-one (D36)

D33 (0.79 g, 2.0 mmol) was dissolved in ethanol (70 mL) and N,N-dimethylformamide (30 mL). 10% palladium on charcoal (0.34 g) was added and the mixture hydrogenated at
40 atmospheric pressure for 20 hours. The mixture was filtered through a plug of celite and the solvent removed *in vacuo* to give the title compound as a grey solid (0.58 g, 96%). MH^+ 303.

Description 37

1-(4-Methyl-3-trifluoromethyl-phenyl)-3-(3-hydroxy-4-methoxy-phenyl)-imidazolidin-2-one (D37)

D34 (0.56 g, 1.2 mmol) was partially dissolved in methanol (70 mL) and N,N-dimethylformamide (4 mL). Ammonium formate (0.41 g, 6.5 mmol) and 10% palladium on charcoal (0.1 g) were added and the mixture heated at reflux for 4 hours and 10 minutes. The mixture was allowed to cool to room temperature, filtered through a plug of celite and concentrated *in vacuo*. The residue was partitioned between dichloromethane (75 mL) and water (75 mL). The organic phase was washed with further water (x2) and brine, dried over anhydrous magnesium sulphate and concentrated *in vacuo* to give the title compound as a pink solid (0.41 g, 92%). MH^+ 367.

Descriptions 38 and 39

1-(2-Chloro-phenyl)-3-(3-hydroxy-4-methoxy-phenyl)-imidazolidin-2-one (D38)
1-(3-Hydroxy-4-methoxy-phenyl)-3-phenyl-imidazolidin-2-one (D39)

A mixture of D35 (0.350 g, 0.86 mmol), ammonium formate (0.25 g, 4 mmol) and 10% palladium on charcoal (0.035 g) in methanol (20 mL) and N,N-dimethylformamide (1 mL) was heated at reflux for one hour. The mixture was allowed to cool to room temperature, filtered through a plug of celite and concentrated *in vacuo*. The residue was partitioned between dichloromethane and water. The organic phase was dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The crude mixture was purified by silica gel chromatography to give the two title compounds D38 and D39. D38 was obtained as a white foam (0.044 g, 16%) MH^+ 319/321 and D39 was obtained as a cream solid (0.092 g, 38%) MH^+ 285.

Description 40

1-[3-(2-Benzyloxy-ethoxy)-4-methoxy-phenyl]-3-(3-fluoro-phenyl)-imidazolidin-2-one (D40)

Benzyl 2-bromoethyl ether (0.15 mL, 0.95 mmol) was added to a mixture of D36 (0.26 g, 0.87 mmol), potassium carbonate (0.15 g, 1.1 mmol) and sodium iodide 0.017 g, 0.1 mmol) in N,N-dimethylformamide (10 mL) and heated at 60°C for 16 hours. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (60 mL) and saturated sodium bicarbonate solution (50 mL). The organic phase was washed with brine (25 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The brown oil was triturated with hexane (x3) to give the title compound as a light brown solid (0.34 g, 91%). MH^+ 437.

Description 41

1-(3-Fluoro-phenyl)-3-[3-(2-hydroxy-ethoxy)-4-methoxy-phenyl]-imidazolidin-2-one (D41)

D40 (0.34 g, 0.79 mmol) was dissolved in ethanol (30 mL) and N,N-dimethylformamide (4 mL). 10% Palladium on charcoal (0.05 g) was added and the mixture hydrogenated at atmospheric pressure for 16 hours. The mixture was filtered through a plug of celite and the solvent removed *in vacuo*. This gave the title compound as a white solid (0.18 g, 66%). MH⁺ 347.

Description 42

Methanesulfonic acid 2-{5-[3-(3-fluoro-phenyl)-2-oxo-imidazolidin-1-yl]-2-methoxy-phenoxy}ethyl ester (D42)

Methanesulfonyl chloride (0.04 mL, 0.53 mmol) was added dropwise at 0°C to a solution of D41 (0.175 g, 0.5 mmol) in dichloromethane (16 mL) and triethylamine (0.11 mL, 0.76 mmol). The mixture was allowed to warm to room temperature and stirred for 1 hour and 15 minutes. The mixture was washed with saturated sodium bicarbonate solution (40 mL) and brine (40 mL). The organic phase was dried over anhydrous magnesium sulphate and concentrated *in vacuo* to give the title compound as a white solid (0.208 g, 97%). MH⁺ 425.

Description 43

3-(3-Fluoro-phenyl)-1-(3-hydroxy-4-methoxy-phenyl)-imidazolidine-2,4-dione (D43)

The title compound was prepared from D23 using the method described in Description 36. MH⁺ 317.

Description 44

5-Nitro-3H-benzoxazol-2-one (D44)

2-Amino-4-nitrophenol (26.1 g, 169 mmol) was dissolved in N,N-dimethylformamide (220 mL) and 1,1'-carbonyldiimidazole (30.2 g, 186 mmol) added. The reaction mixture was heated at 80°C for three hours. The mixture was allowed to cool to room temperature and poured into water (1 Litre) and stirred. The solid was filtered off, washed with water and dried *in vacuo* to give the title compound as a pale yellow solid. MH⁺ 179.

Description 45

5-Nitro-3-(2-piperidin-1-yl-ethyl)-3H-benzoxazol-2-one (D45)

D44 (0.36 g, 2 mmol) was dissolved in N,N-dimethylformamide (10 mL) and sodium hydride (60% dispersion in oil, 0.20 g, 8.3 mmol) added portionwise. The reaction mixture was stirred at room temperature for one hour and a solution of 1-(2-chloroethyl)piperidine monohydrochloride (0.37 g, 2 mmol) in N,N-dimethylformamide (2 mL) added dropwise. The reaction mixture was stirred at room temperature for sixteen hours. The solvent was removed *in vacuo* and the residue partitioned between dichloromethane and saturated sodium bicarbonate solution. The organic phase was dried over anhydrous magnesium sulphate and

concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound (0.35 g, 60 %). MH^+ 292.

Description 46

5 5-Amino-3-(2-piperidin-1-yl-ethyl)-3H-benzoxazol-2-one (D46)

D45 was dissolved in ethanol and 10 % palladium on charcoal added. The reaction mixture was hydrogenated at atmospheric pressure for six hours. The mixture was filtered through a plug of celite and the solvent removed *in vacuo* to give the title compound as an oil (0.29 g, 95%). MH^+ 262.

Description 47

15 5-[2-(3,4-Dichloro-phenylamino)-ethylamino]-3-(2-piperidin-1-yl-ethyl)-3H-benzoxazol-2-one (D47)

The title compound was prepared from D4 and D46 using the method described in Description 24. MH^+ 449/451/453.

Example 1

20 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one (E1)

A mixture of D36 (0.1 g, 0.33 mmol) and 1-(2-chloroethyl)piperidine monohydrochloride (0.67 g, 0.36 mmol) in ethylene glycol dimethyl ether (7 mL), water (3 mL) and saturated potassium carbonate solution (2 mL) was heated at reflux for 5 hours. The mixture was diluted with saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (40 mL). The organic phase was washed with brine (20 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a cream coloured solid (0.072 g, 53%).

1H NMR ($CDCl_3$) δ : 7.50-7.45 (2H, m), 7.31-7.28 (2H, m), 6.85-6.76 (3H, m), 4.19 (2H, t, 6.3 Hz), 3.94 (4H, s), 3.85 (3H, s), 2.84 (2H, t, 6.3 Hz), 2.54 (4H, bs), 1.62 (4H, m), 1.26 (2H, m). MH^+ 414.

Example 2

35 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one (E2)

The title compound was prepared from D36 and 4-(2-chloroethyl)morpholine hydrochloride heating for 10 hours using the method described in Example 1.

1H NMR ($CDCl_3$) δ : 7.57 (1H, d, 2.5 Hz), 7.49-7.46 (2H, m), 7.32-7.27 (2H, m), 6.86 (1H, d, 8.7 Hz), 6.80-6.76 (2H, m), 4.19 (2H, t, 6.0 Hz), 3.94 (4H, s), 3.85 (3H, s), 3.73 (4H, t, 4.6Hz), 2.85 (2H, t, 6.0 Hz), 2.60 (4H, t, 4.6 Hz). MH^+ 416.

Example 3**1-(4-Methyl-3-trifluoromethyl-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one (E3)**

- 5 The title compound was prepared from D37 and 1-(2-chloroethyl)piperidine monohydrochloride heating for 10 hours using the method described in Example 1.
¹H NMR (CDCl₃) δ : 7.77 (1H, d, 2.0 Hz), 7.70 (1H, dd, 8.4 Hz, 2.1Hz), 7.26 (1H, m), 6.85 (2H, s), 4.20 (2H, t, 6.3 Hz), 3.93 (4H, s), 3.84 (3H, s), 2.85 (2H, t, 6.3 Hz), 2.55 (4H, bs), 2.44 (3H, d, 1.1 Hz), 1.62 (4H, m), 1.45 (2H, m). MH⁺ 478.

10

Example 4**1-(4-Methyl-3-trifluoromethyl-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one (E4)**

- 15 The title compound was prepared from D37 and 4-(2-chloroethyl)morpholine hydrochloride heating for 12 hours using the method described in Example 1.
¹H NMR (CDCl₃) δ : 7.76 (1H, d, 2.0Hz), 7.71 (1H, dd, 2.0 Hz, 8.4 Hz), 7.56 (1H, d, 2.4 Hz), 7.26 (1H, m), 6.86 (1H, d, 8.8 Hz), 6.79 (1H, dd, 8.8Hz, 2.4 Hz), 4.19 (2H, t, 6.0 Hz), 3.95 (4H, s), 3.84 (3H, s), 3.73 (4H, t, 4.4 Hz), 2.85 (2H, t, 6.0 Hz), 2.60 (4H,t, 4.4 Hz), 2.44 (3H, d, 0.8 Hz). MH⁺ 480.

20

Example 5**1-(2-Chloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one (E5)**

25

- The title compound was prepared from D38 and 1-(2-chloroethyl)piperidine hydrochloride heating for 3 hours and then stirring at room temperature for 16 hours using the method described in Example 1.
¹H NMR (CDCl₃) δ : 7.63 (1H, d, 2.0 Hz), 7.48-7.43 (2H, m), 7.33 (1H, m), 7.26 (2H, m), 6.84-6.82 (2H, m), 4.23 (2H, t, 6.0 Hz), 4.02-3.91 (4H, m), 3.84 (3H, s), 2.91 (2H, broad s), 2.62 (4H, bs), 1.66 (4H, bs), 1.46 (2H, bs). MH⁺ 430/432

30

Example 6**1-[4-Methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-3-phenyl-imidazolidin-2-one (E6)**

35

- The title compound was prepared from D39 and 1-(2-chloroethyl)piperidine hydrochloride heating for 3 hours and then stirring at room temperature for 16 hours using the method described in Example 1.
- 40 ¹H NMR (CDCl₃) δ : 7.64-7.56 (2H, m), 7.37, (2H, m), 7.09 (1H, t, 7.4 Hz), 6.89-6.84 (2H, m), 4.25 (2H, t, 6.0 Hz), 3.99-3.90 (4H, m), 3.85 (3H, s), 2.93 (2H, t, 6.0 Hz), 2.65 (4H, bs), 1.68 (4H, bm), 1.48 (2H, bm). MH⁺ 396

Example 7**1-[3-(2-Dimethylamino-ethoxy)-4-methoxy-phenyl]-3-(3-fluorophenyl)-imidazolidin-2-one (E7)**

5 Dimethylamine solution (2M) in tetrahydrofuran (0.5 mL, 1 mmol) and D42 (0.1 g, 0.24 mmol) in N,N-dimethylformamide (7 mL) with potassium carbonate (0.065 g, 0.47 mmol) and sodium iodide (0.035 g, 0.23 mmol) were heated in a bomb at 80°C for 16 hours. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (50 mL) and saturated sodium bicarbonate solution (20 mL). The organic phase was washed with water
10 (10 mL) and brine (30 mL), dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as a pale yellow foam (0.074 g, 84%).

¹H NMR (CDCl₃) δ : 7.52-7.46 (2H, m), 7.31-7.29 (2H, m), 6.87-6.77 (3H, m), 4.15 (2H, t, 6.0 Hz), 3.93 (4H, s), 3.85 (3H, s), 2.79 (2H, t, 6.0 Hz), 2.35 (6H, s). MH⁺ 374.

15

Example 8**1-(3-Fluoro-phenyl)-3-{4-methoxy-3[(1S,4S)-2-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-ethoxy]-phenyl}-imidazolidin-2-one (E8)**

20 (1S,4S)-(+)-2-Aza-5-oxabicyclo[2.2.1]heptane hydrochloride (0.064 g, 0.47 mmol) and D17 (0.1 g, 0.24 mmol) in N,N-dimethylformamide (10 mL) with potassium carbonate (0.081 g, 0.59 mmol) and sodium iodide (0.035 g, 0.23 mmol) were heated at 80°C for 16 hours. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (60mL) and saturated sodium bicarbonate solution (15 mL). The organic phase was dried over anhydrous
25 magnesium sulphate and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give the title compound as an orange oil (0.051 g, 50%).

¹H NMR (CDCl₃) δ : 7.54 (1H, d, 2.4 Hz), 7.49-7.46 (1H, m), 7.32-7.29 (2H, m), 6.86 (1H, d, 8.8 Hz), 6.82-6.76 (2H, m), 4.40 (1H, s), 4.15 (2H, t, 6.4 Hz), 4.08 (1H, d, 8.0 Hz), 3.94 (4H, s), 3.85 (3H, s), 3.66-3.62 (2H, m), 3.10-3.02 (3H, m), 2.64 (1H, d, 10 Hz), 1.88 (1H, d, 10
30 Hz), 1.74 (1H, d, 10 Hz). MH⁺ 428.

Example 9**1-(3,4-Dichloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one (E9)**

35

Phosgene solution (~20% in toluene, 0.34 mL, 0.7 mmol) was added dropwise at room temperature to a solution of D27 (0.30 g, 0.7 mmol) in tetrahydrofuran (15 mL) and triethylamine (0.34 mL, 2.4 mmol) and stirred at room temperature for 16 hours. Ethyl acetate (120 mL) was added and the reaction mixture washed with 2M sodium hydroxide
40 solution (x3) and brine (60 mL). The organic phase was dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The crude compound was purified by silica gel chromatography to give the title compound as a beige solid (0.219 g, 69%).

¹H NMR (CDCl₃) δ : 7.72 (1H, d, 2.5 Hz), 7.50 (1H, dd, 8.8 Hz, 2.5 Hz), 7.47 (1H, d, 2.0 Hz), 7.40 (1H, d, 9.1 Hz), 6.87-6.84 (2H, m), 4.18 (2H, t, 6.3 Hz), 3.97-3.87 (4H, m), 3.85 (3H, s), 2.83 (2H, t, 6.3 Hz), 2.52 (4H, m), 1.61 (4H, m), 1.45 (2H, m). MH⁺ 464/466/468.

5 **Example 10**

1-(3,4-Dichloro-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one (E10)

The title compound was prepared from D28 using the method described in Example 9

10 ¹H NMR (CDCl₃) δ : 7.72 (1H, d, 2.4 Hz), 7.54-7.48 (2H, m), 7.40 (1H, d, 8.8 Hz), 6.86 (1H, d, 8.8 Hz), 6.80 (1H, dd, 2.4 Hz), 4.19 (2H, t, 6.0 Hz), 3.97-3.90 (4H, m), 3.85 (3H, s), 3.73 (4H, t, 4.8 Hz), 2.85 (2H, t, 6.0 Hz), 2.60 (4H, t, 4.8 Hz). MH⁺ 466/468/470.

Example 11

15 **1-(3,4-Dichloro-phenyl)-3-[3-(2-dimethylamino-ethoxy)-4-methoxy-phenyl]-imidazolidin-2-one (E11)**

The title compound was prepared from D29 using the method described in Example 9.

20 ¹H NMR (CDCl₃) δ : 7.72 (1H, d, 2.4 Hz), 7.50-7.47 (2H, m), 7.38 (1H, d, 8.8 Hz), 6.84 (2H, m), 4.15 (2H, t, 6.0 Hz), 3.95-3.88 (4H, m), 3.84 (3H, s), 2.79 (2H, t, 6.0 Hz), 2.35 (6H, s). MH⁺ 424/426/428.

Example 12

25 **1-(3,4-Dichloro-phenyl)-3-[6-(2-piperidin-1-yl-ethoxy)-pyridin-3-yl]-imidazolidin-2-one (E12)**

The title compound was prepared from D30 using the method described in Example 9.

30 ¹H NMR (CDCl₃) δ : 8.10-8.06 (2H, m), 7.73 (1H, d, 2.8 Hz), 7.50 (1H, dd, 8.8 Hz, 2.8 Hz), 7.40 (1H, d, 8.8 Hz), 6.81-6.78 (1H, m), 4.43 (2H, t, 6.0 Hz), 3.96 (4H, m), 2.76 (2H, t, 6.0 Hz), 2.51 (4H, m), 1.60 (4H, m), 1.44 (2H, m). MH⁺ 435/437/439.

Example 13

35 **1-[4-Bromo-3-(2-piperidin-1-yl-ethoxy)-phenyl]-3-(3,4-dichloro-phenyl)-imidazolidin-2-one (E13)**

The title compound was prepared from D31 using the method described in Example 9.

40 ¹H NMR (CDCl₃) δ : 7.72 (1H, d, 2.5 Hz), 7.59 (1H, d, 2.5 Hz), 7.51-7.39 (3H, m), 6.77 (1H, dd, 8.7 Hz, 2.5 Hz), 4.17 (2H, m), 3.90 (4H, m), 2.86 (2H, m), 2.58 (4H, m), 1.62 (4H, m), 1.44 (2H, m). MH⁺ 512/514/516/518.

Example 14

1-(2,3-Dichloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one (E14)

The title compound was prepared from D32 using the method described in Example 9.

- 5 ¹H NMR (CDCl₃) δ : 7.61 (1H, d, 2.4 Hz), 7.44 (1H, dd, 8.0 Hz, 1.6 Hz), 7.36 (1H, dd, 1.6 Hz, 8.0 Hz), 7.27 (1H, m), 6.85 (1H, 8.8 Hz), 6.80 (1H, m), 4.16 (2H, t, 6.4Hz), 4.00 (2H, m), 3.90 (2H, m), 3.85 (3H, s), 2.82 (2H, 6.4 Hz), 2.50 (4H, m), 1.58 (4H, m), 1.43 (2H, m). MH⁺ 464/466/468.

10 **Example 15**

1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-1,3-dihydro-imidazol-2-one (E15)

- 15 D20 (0.110 g, 0.23 mmol) in toluene (12 mL) was treated with concentrated hydrochloric acid (one drop) and the mixture heated at reflux for three hours. The mixture was diluted with ethyl acetate and washed with saturated sodium bicarbonate solution (25 mL) and brine (40 mL). The organic phase was dried over anhydrous magnesium sulphate and concentrated *in vacuo*. The crude compound was purified by silica gel chromatography to give the title compound (0.049 g, 52%).

- 20 ¹H NMR (CDCl₃) δ : 7.52-7.49 (1H, m), 7.42-7.39 (2H, m), 7.30 (1H, d, 2.4 Hz), 7.05 (1H, dd, 2.4 Hz, 8.4 Hz), 6.96 (1H, m), 6.91 (1H, d, 8.4 Hz), 6.71 (2H, dd, 12.8 Hz, 3.2 Hz), 4.19 (2H, t, 6.4 Hz), 3.88 (3H, s), 2.84 (2H, t, 6.4 Hz), 2.52 (4H, m), 1.60 (4H, m), 1.45 (2H, m). MH⁺ 412.

25 **Example 16**

1-(3,5-Difluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-1,3-dihydro-imidazol-2-one (E16)

The title compound was prepared from D21 using the method described in Example 15.

- 30 ¹H NMR (CDCl₃) δ : 7.34-7.30 (2H, m), 7.25 (1H, m), 7.03 (1H, dd, 8.6 Hz, 2.5 Hz), 6.92 (1H, d, 8.6 Hz), 6.73-6.68 (3H, m), 4.18 (2H, t, 6.4 Hz), 3.88 (3H, s), 2.83 (2H, t, 6.4 Hz), 2.52 (4H, m), 1.59 (4H, m), 1.44 (2H, m). MH⁺ 430.

Example 17

- 35 **1-(3,5-Difluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazol-2-one (E17)**

E16 (0.064 g, 0.15 mmol) was dissolved in ethanol (15 mL) and 10% palladium on charcoal added (0.02 g). The reaction mixture was hydrogenated at atmospheric pressure for 48 hours.

- 40 The mixture was filtered through a small plug of celite and the solvent removed *in vacuo*. The residue was triturated with petroleum ether (40-60°C) (x3) to give the title compound (0.035 g, 55%).

¹H NMR (CDCl₃) δ : 7.45 (1H, d, 1.7 Hz), 7.22-7.17 (2H, m), 6.88-6.85 (2H, m), 6.52 (1H, m), 4.18 (2H, t, 6.4 Hz), 3.97-3.88 (4H, m), 3.85 (3H, s), 2.83 (2H, t, 6.4 Hz), 2.51 (4H, m), 1.62 (4H, m), 1.46 (2H, m). MH⁺ 432.

5 **Example 18**

3-(3-Fluoro-phenyl)-1-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazol-2,4-dione (E18)

10 Diethyl azodicarboxylate (0.055 g, 0.32 mmol) in tetrahydrofuran (1mL) was added dropwise to a stirred solution of D43 (0.1 g, 0.32 mmol), triphenylphosphine (0.083 g, 0.32 mmol) and N-(2-hydroxyethyl)piperidine (0.041 g, 0.32 mmol) in tetrahydrofuran (4 mL) at room temperature. After stirring for 24 hours the reaction mixture was passed through an ion exchange SCX cartridge eluting with methanol and 0.5 M ammonia/methanol. The ammonia fractions were combined and concentrated *in vacuo*. The residue was purified by silica gel chromatography followed by trituration with petroleum ether (40-60°C) (x2) to give the title compound as a white solid. (0.062 g, 54%).

15 ¹H NMR (CDCl₃) δ : 7.50-7.45 (2H, m), 7.32-7.24 (2H, m), 7.15-7.02 (1H, m), 6.87 (2H, m), 4.45 (2H, s), 4.17 (2H, t, 6.4 Hz), 3.87 (3H, s), 2.83 (2H, t, 6.4 Hz), 2.51 (4H, m), 1.60 (4H, m), 1.45 (2H, m). MH⁺ 428

20

Example 19


5-[3-(3,4-Dichloro-phenyl)-2-oxo-imidazolidin-1-yl]-3-(2-piperidin-1-yl-ethyl)-3H-benzooxazol-2-one (E19)

25 The title compound was prepared from D47 using the method described in Example 9

¹H NMR (CDCl₃) δ : 7.73 (1H, d, 2.4 Hz), 7.71 (1H, d, 2.4 Hz), 7.50 (1H, m), 7.42 (1H, d, 8.8 Hz), 7.16 (1H, d, 8.8 Hz), 6.92 (1H, dd, 8.8 Hz, 2.4 Hz), 4.04-3.93 (6H, m), 2.67 (2H, t, 6.8 Hz), 2.47 (4H, bm), 1.54 (4H, bm), 1.41 (2H, bm). MH⁺ 475/477/479.

Claims

1. A compound of formula (I):



(I)

5 wherein:

a is 0, 1, 2, 3, 4 or 5;

b is 1, 2 or 3;

Y is nitrogen or carbon;

A is oxygen, nitrogen, -CONH-, -NHCO- or together with R² form a benzoxazolone group;

10 R¹ is halogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, hydroxy, amino, mono- or di-C₁₋₆alkylamino, nitro, CN, CF₃, OCF₃, aryl, arylC₁₋₆alkyl, arylC₁₋₆alkyloxy or arylC₁₋₆alkylthio;

R² is hydrogen, halogen, C₁₋₆alkyl, C₁₋₆alkoxy, haloC₁₋₆alkyl or haloC₁₋₆alkoxy;

R³ is:

- 15 (i) -NR⁴R⁵ where R⁴ and R⁵ are independently hydrogen, C₁₋₆alkyl or arylC₁₋₆alkyl; or
 (ii) an optionally substituted N-linked heterocycle; or
 (iii) an optionally substituted C-linked heterocycle;

20 ----- is a single bond or a double bond;

X is CH₂ or C=O (when ----- is a single bond) or X is CH (when ----- is a double bond).

2. A compound as claimed in claim 1, wherein a is 1 or 2.


3. A compound as claimed in claim 1 or claim 2, wherein R¹ is halogen.

25 4. A compound as claimed in claim 1, 2 or 3, wherein R¹ is at the 3 position or the 4 position.


5. A compound as claimed in any of claims 1 to 4, wherein X is CH₂ and ----- represents a single bond or X is CH and ----- represents a double bond.

6. A compound as claimed in any of claims 1 to 6, wherein Y is carbon.

30 7. A compound as claimed in any of claims 1 to 6, wherein R² is at the following position:




8. A compound as claimed in any of claims 1 to 7, wherein R^2 is C_{1-6} alkoxy.
9. A compound as claimed in any of claims 1 to 8, wherein A is at the following position:



- 5
10. A compound as claimed in any of claims 1 to 9, wherein A is oxygen.
11. A compound as claimed in any of claims 1 to 10, wherein b is 2.
12. A compound as claimed in any of claims 1 to 11, wherein R^3 is N-linked heterocycle or di- C_{1-6} alkylamine.
- 10 13. A compound as claimed in claim 1, which is:
- 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one;
- 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one;
- 1-(4-Methyl-3-trifluoromethyl-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one;
- 15 1-(4-Methyl-3-trifluoromethyl-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-one;
- 1-(2-Chloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one
- 1-[4-Methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-3-phenyl-imidazolidin-2-one;
- 1-[3-(2-Dimethylamino-ethoxy)-4-methoxy-phenyl]-3-(3-fluorophenyl)-imidazolidin-2-one;
- 20 1-(3-Fluoro-phenyl)-3-{4-methoxy-3[(1S,4S)-2-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-ethoxy]-phenyl}-imidazolidin-2-one;
- 1-(3,4-Dichloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one;
- 1-(3,4-Dichloro-phenyl)-3-[4-methoxy-3-(2-morpholin-4-yl-ethoxy)-phenyl]-imidazolidin-2-
- 25 one;
- 1-(3,4-Dichloro-phenyl)-3-[3-(2-dimethylamino-ethoxy)-4-methoxy-phenyl]-imidazolidin-2-one;
- 1-(3,4-Dichloro-phenyl)-3-[6-(2-piperidin-1-yl-ethoxy)-pyridin-3-yl]-imidazolidin-2-one;
- 1-[4-Bromo-3-(2-piperidin-1-yl-ethoxy)-phenyl]-3-(3,4-dichloro-phenyl)-imidazolidin-2-one;
- 30 1-(2,3-Dichloro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazolidin-2-one;
- 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-1,3-dihydro-imidazol-2-one;
- 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-1,3-dihydro-imidazol-2-one;
- 35 1-(3-Fluoro-phenyl)-3-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazol-2-one;
- 3-(3-Fluoro-phenyl)-1-[4-methoxy-3-(2-piperidin-1-yl-ethoxy)-phenyl]-imidazol-2,4-dione;

5-[3-(3,4-Dichloro-phenyl)-2-oxo-imidazolidin-1-yl]-3-(2-piperidin-1-yl-ethyl)-3H-benzoxazol-2-one,
or a pharmaceutically acceptable salt thereof.


- 5 14. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises:
a) the cyclisation of a compound of formula (II)



(II)


10

in which R¹, R², R³, A, Y, a and b are defined in formula (I); or
b) the coupling of a compound of formula (III):




(III)

- 15 in which R¹, R², A, X, Y, a and b are defined in formula (I) and L¹ is a leaving group, with a compound of formula (IV):



(IV)


in which R⁴ and R⁵ are defined in formula (I); or
c) the coupling of a compound of formula (V)



(V)

20

in which R¹, R², X, Y and a are defined in formula (I), with a compound of formula (VI):



(VI)

- 25 in which R⁴, R⁵ and b are defined in formula (I) and L² is a leaving group or hydroxy; or
d) the cyclisation of a compound of formula (VII):

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/GB 03/00020

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4453 A61K31/5355 A61K31/4166 A61K31/423 A61K31/44
C07D233/70 C07D233/78 C07D413/04 C07D401/04 C07D491/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 23783 A (DAVIES DAVID THOMAS ; GASTER LARAMIE MARY (GB); WYMAN PAUL ADRIAN () 8 August 1996 (1996-08-08) cited in the application page 1, line 8 - line 10; claim 1	1-23
A	WO 97 48700 A (BROMIDGE STEVEN MARK ; SMITHKLINE BEECHAM PLC (GB)) 24 December 1997 (1997-12-24) cited in the application page 1, line 8 - line 11	1-23



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Δ* document member of the same patent family

Date of the actual completion of the international search

28 March 2003

Date of mailing of the international search report

03/04/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 03/00020

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9623783	A	08-08-1996	AP 657 A	06-08-1998
			AT 197300 T	15-11-2000
			AU 699727 B2	10-12-1998
			AU 4664696 A	21-08-1996
			BG 101806 A	30-04-1998
			BR 9607016 A	28-10-1997
			CA 2212061 A1	08-08-1996
			CZ 9702445 A3	16-09-1998
			DE 69610822 D1	07-12-2000
			DE 69610822 T2	07-06-2001
			DK 808312 T3	12-02-2001
			EA 304 B1	29-04-1999
			WO 9623783 A1	08-08-1996
			EP 0808312 A1	26-11-1997
			ES 2151652 T3	01-01-2001
			FI 973205 A	01-10-1997
			GR 3035075 T3	30-03-2001
			HK 1003883 A1	31-08-2001
			HU 9901115 A2	28-07-1999
			IL 116998 A	08-08-2001
			JP 10513442 T	22-12-1998
			NO 973543 A	01-10-1997
			NZ 301265 A	23-12-1998
			PL 321706 A1	22-12-1997
			PT 808312 T	30-03-2001
			RO 115522 B	30-03-2000
			SI 808312 T1	28-02-2001
			SK 103897 A3	04-02-1998
			TR 9700749 T1	21-02-1998
			US 6235758 B1	22-05-2001
			US 5990133 A	23-11-1999
			ZA 9600758 D	30-09-1997
WO 9748700	A	24-12-1997	AT 196766 T	15-10-2000
			AU 718000 B2	06-04-2000
			AU 3339697 A	07-01-1998
			BR 9709982 A	10-08-1999
			CA 2258559 A1	24-12-1997
			CZ 9804204 A3	12-05-1999
			DE 69703242 D1	09-11-2000
			DE 69703242 T2	08-03-2001
			DK 912556 T3	22-01-2001
			WO 9748700 A1	24-12-1997
			EP 0912556 A1	06-05-1999
			ES 2152682 T3	01-02-2001
			GR 3034847 T3	28-02-2001
			HK 1019744 A1	15-03-2002
			HU 9903480 A2	28-07-2000
			JP 2000512644 T	26-09-2000
			KR 2000022044 A	25-04-2000
			NO 985970 A	18-12-1998
			NZ 332840 A	26-05-2000
			PL 330708 A1	24-05-1999
			PT 912556 T	28-02-2001
			SI 912556 T1	28-02-2001
			TR 9802552 T2	22-02-1999
			US 6313145 B1	06-11-2001
			US 2002035134 A1	21-03-2002

Information on patent family members

PCT/GB 03/00020

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9748700 A		ZA 9705416 A	25-01-1999